

- High energy level
- High attractivity
- Good performance
- Semi-intensive farming
- Good for autumn feeding



COMPOSITION:

Analyses (%)		Sizes
Protein	34	3.0 mm
Fat	15	4.5 mm
Crude fibre	1.5	6.0 mm
Ash	5.0	8.0 mm
Total P	0.97	14.0 mm
		20.0 mm

Vitamins added

Vitamin A (IE/kg)	10000
-------------------	-------

Energy (MJ/kg)

Gross Energy	20.6
Digestible Energy	17.4

FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR):


Fish weight (g)	Feed size (mm)	< 10 °C	10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C	> 26 °C
50-100	3.0	According to fish's appetite	1.00	1.50	2.00	2.61	3.11	3.61	4.11	4.61	4.06	According to fish's appetite and O2 level
100-250	4.5		0.80	1.00	1.50	1.80	2.31	2.81	3.31	3.81	3.35	
250-500	6.0		0.60	0.80	1.00	1.30	1.50	1.80	2.31	2.81	2.47	
500-1000	6.0		0.50	0.60	0.80	1.00	1.30	1.50	1.80	2.00	1.76	
1000-2000	8.0		0.40	0.50	0.70	0.80	1.10	1.30	1.60	1.80	1.59	

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.